

Table 1 The decrease (-) or increase in the tobacco use in different states in India from 2009-10 to 2016-17 (Global Adult Survey)

State	2009-10	2016-17	% Change
Sikkim	41.6	17.9	-56.97
Bihar	53.5	25.9	-51.59
Kerala	21.4	12.7	-40.65
Andhra Pradesh	29.2	20	-31.51
Delhi	24.3	17.8	-26.75
Chhattisgarh	53.2	39.1	-26.50
Puducherry	15.1	11.2	-25.83
Himachal Pradesh	21.2	16.1	-24.06
Nagaland	56.8	43.3	-23.77
Rajsthan	32.3	24.7	-23.53
Jharkhand	50.1	38.9	-22.36
Karnatka	28.2	22.8	-19.15
India	34.6	28.6	-17.34
Maharashtra	31.4	26.6	-15.29
Meghalya	55.2	47	-14.86
Gujrat	29.4	25.1	-14.63
Utrakhand	30.7	26.5	-13.68
Madhya Pradesh	39.5	34.2	-13.42
Mijoram	67.2	58.7	-12.65
Jammu & Kashmir	26.6	23.7	-10.90
West Bengal	36.3	33.5	-7.71
Arunachal Pradesh	47.7	45.5	-4.61
Chandigarh	14.3	13.7	-4.20
Orissa	46.2	45.6	-1.30
Haryana	23.7	23.6	-0.42
Manipur	54.1	55.1	1.85
Uttar Pradesh	33.9	35.5	4.72
Goa	8.8	9.7	10.23
Punjab	11.7	13.4	14.53
Tripura	55.9	64.5	15.38
Asam	39.3	48.2	22.65
Tamilnadu	16.2	20	23.46